

Low energy processes to distinguish among seesaw models

C. BIGGIO

Max-Planck-Institut für Physik, 80805 München, Germany

Summary. — We consider the three basic seesaw scenarios (with fermionic singlets, scalar triplets or fermionic triplets) and discuss their phenomenology, aside from neutrino masses. We use the effective field theory approach and compare the dimension-six operators characteristic of these models. We discuss the possibility of having large dimension-six operators and small dimension-five (small neutrino masses) without any fine-tuning, if the lepton number is violated at a low energy scale. Finally, we discuss some peculiarities of the phenomenology of the fermionic triplet seesaw model.

PACS 13.15.+g 14.60.Pq 14.60.St – .

Neutrinos are exactly massless in the Standard Model (SM). However, in the past decades, oscillation experiments have proved that neutrinos are massive and nowadays we know they are lighter than few eV, thanks to 3H decay experiments, $0\nu\beta\beta$ decay experiments and, more recently, to cosmology.

New physics (NP) is therefore required to explain neutrino masses and a natural explanation of their smallness, with respect to charged leptons ones, can be achieved in the context of seesaw models, which introduce NP at an energy scale higher than the electroweak (EW) one. A good way of considering the effects of this new high energy physics is using the effective field theory approach. While with SM fields many dimension-six ($d6$) and higher order operators can be built, the $d5$ is unique and it is given by $\left[\frac{1}{2}c_{\alpha\beta}^{d5}\left(\bar{\ell}_{L\alpha}^c\tilde{\phi}^*\right)\left(\tilde{\phi}^\dagger\ell_{L\beta}\right)+\text{h.c.}\right]$, where ℓ_L are the lepton doublets and $\tilde{\phi} = i\tau_2\phi^*$, with ϕ the Higgs field. Here $c_{\alpha\beta}^{d5}$ is a model-dependent coefficient matrix inversely proportional to the NP scale M . After EW symmetry breaking this operator generates Majorana masses for neutrinos which, for large enough M , are small for “natural” values of the Yukawa couplings ($Y \sim O(10^{-6} - 1)$).

The unique $d5$ operator can be generated by different types of NP. At tree-level, there are only three possibilities: from heavy fermionic singlets (type-I seesaw), scalar triplets (type-II seesaw) or fermionic triplets (type-III seesaw). It is clear that from neutrino masses it is impossible to understand which is the NP which lays behind, so that, in order to distinguish among different models, other associated effects have to be observed. $d6$ operators, which produce them, have been computed in refs. [1, 2], respectively for the type-I model and types II and III, and are shown in table I, together with the

TABLE I. – *Coefficients of the $d5$ operator, c^{d5} , and $d6$ operators and their coefficients, c^{d6} , in the three basic seesaw theories.*

Model	c^{d5}	c_i^{d6}	$\mathcal{O}_i^{d6}$
Fermionic Singlet	$Y_N^T \frac{1}{M_N} Y_N$	$\left(Y_N^\dagger \frac{1}{M_N^\dagger} \frac{1}{M_N} Y_N \right)_{\alpha\beta}$	$\left(\overline{\ell_{L\alpha}} \widetilde{\phi} \right) i \not{\partial} \left(\widetilde{\phi}^\dagger \ell_{L\beta} \right)$
Scalar Triplet	$4Y_\Delta \frac{\mu_\Delta}{M_\Delta^2}$	$\frac{1}{M_\Delta^2} Y_{\Delta\alpha\beta} Y_{\Delta\gamma\delta}^\dagger$	$\left(\overline{\ell_{L\alpha}} \overline{\tau} \ell_{L\beta} \right) \left(\overline{\ell_{L\gamma}} \overline{\tau} \ell_{L\delta} \right)$
		$\frac{ \mu_\Delta ^2}{M_\Delta^4}$ $-2(\lambda_3 + \lambda_5) \frac{ \mu_\Delta ^2}{M_\Delta^4}$	$\left(\phi^\dagger \overline{\tau} \widetilde{\phi} \right) \left(\overline{D_\mu} \overline{D^\mu} \right) \left(\widetilde{\phi}^\dagger \overline{\tau} \phi \right)$ $\left(\phi^\dagger \phi \right)^3$
Fermionic Triplet	$Y_\Sigma^T \frac{1}{M_\Sigma} Y_\Sigma$	$\left(Y_\Sigma^\dagger \frac{1}{M_\Sigma^\dagger} \frac{1}{M_\Sigma} Y_\Sigma \right)_{\alpha\beta}$	$\left(\overline{\ell_{L\alpha}} \overline{\tau} \widetilde{\phi} \right) i \not{\partial} \left(\widetilde{\phi}^\dagger \overline{\tau} \ell_{L\beta} \right)$

corresponding coefficients (and the coefficients of the $d5$ operators). We immediately observe that the $d6$ operators differ in the three models, so that in principle, observing the physical effects they produce, it would be possible to understand which is that one responsible for neutrino masses if among these three.

Before starting the analysis of the phenomenology associated to $d6$ operators, we discuss here about the possibility of observing it. If we consider $\mathcal{O}(1)$ Yukawa couplings and we look at the fermionic seesaws, we see that $c^{d6} \sim (c^{d5})^2$. Since c^{d5} corresponds to neutrino masses, it must be small, implying that the effects coming from the $d6$ operators are negligible. However, if we look at the type-II model, we observe that we can have a relatively large c^{d6} (here we are referring to the first of the list, which is the phenomenologically relevant one) with large Yukawa couplings and small neutrino masses if the scale μ_Δ is small enough. For example, if $Y \sim \mathcal{O}(1)$ and $M \sim 1\text{TeV}$, then $\mu_\Delta \sim 1\text{eV}$ would give the correct order of magnitude for neutrino masses. In this model μ_Δ is the dimension-full trilinear coupling among two Higgs doublets and the new scalar triplet and it is the scale associated to lepton number violation. Since $d6$ operators are lepton number conserving, we can suppose [2] that whenever the lepton number breaking scale μ is small and does not coincide with the scale of NP M , neutrino masses will be directly proportional to μ , while c^{d6} will remain insensitive to this:

$$(1) \quad c^{d5} = f(Y) \frac{\mu}{M^2} \quad c^{d6} = g(Y) \frac{1}{|M|^2},$$

where f and g are some functions of Yukawa couplings. If M is low enough, effects coming from NP responsible for neutrino masses can be observed without invoking any fine-tuning, since the $d5$ and $d6$ operators are now decoupled. As already mentioned, this is “naturally” realized in the type-II seesaw model. As for fermionic seesaws, particular textures of heavy Majorana masses have to be chosen [3], but it is still feasible.

Once established that the seesaw scale can be lowered without introducing any fine-tuning to maintain neutrino masses small, the associated effects produced by different $d6$ operators can be analyzed. The kinetic corrections given by the $d6$ operator in the type-I and type-III models produce a non-unitary mixing matrix in the SM charged currents and flavor changing neutral currents (FCNC) in the neutrino sector [1, 4]. Moreover, only in

the type-III case, FCNC are present in the charged lepton sector too [2]. Among the three $d6$ operators characteristic of the type-II model, only the first one is phenomenologically relevant, since the others only renormalize the SM parameters: this operator generates flavor-changing four-fermion interactions that can be easily tested. In ref. [2] a detailed analysis of EW decays and flavor-changing processes in the three models have been carried out and bounds on c^{d6} have been derived. The general trend is the following: for $M \sim 1$ TeV, $Y < 10^{-1}$ or smaller.

Before concluding we discuss briefly some peculiarities of the phenomenology of the type-III seesaw model. The presence of FCNC in the charged lepton sector allows processes like $\mu \rightarrow eee$ and flavor violating τ decays already at tree-level. This permits to derive strong bounds on the off-diagonal elements of the matrix $c_{\alpha\beta}^{d6}$, stronger than the ones that can be derived from radiative processes. However, in ref. [5] the $\mu \rightarrow e\gamma$ and $\tau \rightarrow l\gamma$ decays have been calculated and the following relationship among branching ratios has been derived:

$$(2) \quad Br(\mu \rightarrow e\gamma) = 1.3 \cdot 10^{-3} \cdot Br(\mu \rightarrow eee)$$

and similarly for τ decays. This relationship is very interesting because, as far as we know, this is the only model in which radiative decays are so much suppressed with respect to non-radiative ones. Moreover, since the current bound on $Br(\mu \rightarrow eee)$ is 10^{-12} and the reach of future experiments on $\mu \rightarrow e\gamma$ is 10^{-14} , a positive signal from these experiments would rule out this model as the unique source of lepton flavor violation.

Additionally in ref. [6] the anomalous magnetic moment of leptons has been calculated and it has been shown that, even if the scale M is pushed down to the EW scale, the contribution of the type-III seesaw is still smaller than (or comparable with) the SM error on this observable, rendering thus impossible to measure it. Moreover it turns out that this contribution has the “wrong” sign in order to explain the measured discrepancy in the muon sector.

To summarize, we have shown that the physical processes associated to the $d6$ operators coming from different seesaw models can be used to distinguish among them, if the scale of new physics is low enough. We have shown that this is possible if neutrino masses suppression is due to a small scale at which lepton number breaking is realized, different from the scale of the new heavy particle. Finally some details on the phenomenology of the type-III seesaw model have been discussed.

REFERENCES

- [1] A. Broncano, M. B. Gavela and E. Jenkins, Phys. Lett. B **552** (2003) 177 [Erratum-ibid. B **636** (2006) 330] [arXiv:hep-ph/0210271]; A. De Gouvea, G. F. Giudice, A. Strumia and K. Tobe, Nucl. Phys. B **623** (2002) 395 [arXiv:hep-ph/0107156].
- [2] A. Abada, C. Biggio, F. Bonnet, M. B. Gavela and T. Hambye, JHEP **0712** (2007) 061 [arXiv:0707.4058 [hep-ph]].
- [3] M. C. Gonzalez-Garcia and J. W. F. Valle, Phys. Lett. B **216** (1989) 360; J. Kersten and A. Y. Smirnov, Phys. Rev. D **76** (2007) 073005 [arXiv:0705.3221 [hep-ph]].
- [4] S. Antusch, C. Biggio, E. Fernandez-Martinez, M. B. Gavela and J. Lopez-Pavon, JHEP **0610** (2006) 084 [arXiv:hep-ph/0607020].
- [5] A. Abada, C. Biggio, F. Bonnet, M. B. Gavela and T. Hambye, arXiv:0803.0481 [hep-ph].
- [6] C. Biggio, arXiv:0806.2558 [hep-ph].